

Classification, Maturity, Provenance, Tectonic Setting and Source-Area Weathering of the Ilubirin Stream Sediments, South West, Nigeria

Henry Y. Madukwe¹✉, Romanus A. Obasi¹

¹Department of Geology, Ekiti State University, Ado-Ekiti, Ekiti State, Nigeria

Abstract: Geochemical study on the Ilubirin stream sediments was carried out to infer their provenance, maturity, classification, enrichment, depletion, tectonic setting and source-area weathering. The sediments can be classified as ferromagnesian potassic, quartz arenites that are non-calcareous. The $\text{SiO}_2/\text{Al}_2\text{O}_3$ index is high, meaning that the samples are mature, the Index of Compositional Variability also indicate that the sediments are mineralogically mature, while the $\text{Al}_2\text{O}_3/(\text{CaO}+\text{MgO}+\text{Na}_2\text{O}+\text{K}_2\text{O})$ ratio indicate that there are stable mobile oxides in the sediments. The plot of SiO_2 versus $\text{Al}_2\text{O}_3 + \text{K}_2\text{O} + \text{Na}_2\text{O}$ shows that the sediments formed under semi-arid/arid conditions tending towards increasing chemical maturity. Al_2O_3 correlates positively with all the major oxides except SiO_2 , suggesting hydraulic fractionation and sorting. The negative linear trend between Al_2O_3 and SiO_2 indicates that the major element composition of the stream sediments is controlled largely by the relative amount of quartz and feldspar versus clay minerals. The depletion of highly mobile Na, K and Ca elements is due to leaching during the formation of clay minerals during increased chemical weathering. The immobile Fe and the less mobile Mg elements were depleted while the immobile Ti was enriched; this suggest that they may be from a felsic source. The high $\text{TiO}_2/\text{Fe}_2\text{O}_3$ ratios suggests concentration in the sediments of a heavy mineral phase containing Ti minerals such as ilmenite and rutile. The weathering indices (CIA, CIW, PIA and MIA) indicates a high degree of weathering of the source materials. The tectonic setting is the passive continental margin, while the provenance is the quartz-rich sediments of mature continental provenance, associated with a continental passive margin, intracratonic basins, or recycled orogenic provinces.

Keywords: weathering, chemical maturity, provenance, felsic, tectonic setting

1.0 Introduction

The geochemical compositions of stream sediments reflect the average composition of an entire drainage basin (Halamic et al., 2001; Reimann and Melezhik, 2001). According to (Grunsky and Sutphin, 2009), geochemical studies based on the chemical analysis of active stream sediments are an effective tool with several applications. Several authors have used major element discrimination diagrams (Bhatia, 1983) to discriminate the tectonic settings of sedimentary basins and have been applied in topical publications (Kroonenberg, 1994; Zimmermann and Bahlburg, 2003; Armstrong-Altrin et al., 2004). According to (Armstrong-Altrin and Verma, 2005), caution is required in their uncritical use. The most important

clues for the tectonic setting of the basin comes from the relative depletion of the most mobile elements like Ca and Na and enrichment of Si and Ti—the most immobile elements, among others. The oxides of these elements are assumed to show enrichment or depletion in quartz, K-feldspar, plagioclase feldspars, and micas. The ratio of the most immobile elements to the most mobile ones increases toward passive margins due to the relative tectonic stability (Bhatia, 1983; Kroonenberg, 1994; Zimmermann and Bahlburg, 2003; Armstrong-Altrin et al., 2004; Roser and Korsch, 1986) and hence prolonged weathering. This can be recorded in sediments as paleoclimate index (Chittleborough, 1991; Harnois, 1988; Nesbitt and Young, 1982) and high degree of sediment recycling.

This article is published under the terms of the Creative Commons Attribution License 4.0

Author(s) retain the copyright of this article. Publication rights with Alkhaer Publications.

Published at: <http://www.ijsciences.com/pub/issue/2015-04/>

Article Number: V420150404; Online ISSN: 2305-3925; Print ISSN: 2410-4477



Henry Y. Madukwe (Correspondence)



henrymadukwe@yahoo.co.uk



+2348063786674

Major elements and selected trace and rare earth elements and their elemental ratios are sensitive indicators of the source rocks, tectonic setting, paleoweathering conditions and paleoclimate of the clastic sediments (Bhatia, 1983; Bhatia and Crook, 1986; Roser and Korsch, 1986; Roser and Korsch, 1988; McLennan and Taylor, 1991; McLennan et al., 1993; Johnsson and Basu, 1993; Condie, 1993; Nesbitt, 1996; Fedo, et al., 1997; Cullers and Podkovyrov, 2000; Bhatt and Ghosh, 2001). Ilubirin stream is located in Ondo State South Western Nigeria.

The study area lies within the latitudes $6^{\circ} 28'N$ and $6^{\circ} 37'N$ and longitudes $4^{\circ} 32'E$ and $4^{\circ} 5'E$ of the Greenwich Meridian (Fig 1). The Elevation ranges between 50 and 250 m above the sea level. This area lies between Ekiti State and Edo State respectively. The drainage pattern is mainly Dendritic in which there are many rivers of different sizes. Dendritic system forms in V-shape valleys as a result of the rock types whether porous or non-porous. The drainage in the area is influenced by: lithological variations, structural elements such as faults, joints etc., and geomorphology of the area. The most outstanding characteristics of the drainage systems over the areas of Basement Complex rocks is the proliferation of many small rivers, some of which intersect the minor roads and footpaths in the area. The area has little streams which take their source from hills in the northern part. This present research is aimed at interpreting the sediment source area weathering, provenance, tectonic setting, maturity and classification of the Ilubirin stream sediments based on major oxides geochemical data.

2.0 Materials and Methods

Seven stream sediments were collected at Ilubirin; the samples were collected using a hand auger and shovel. They were then sieved after drying using $75\mu m$ stainless steel mesh wire. Chemical analysis was carried at Stellenbosch University, South Africa

using the Inductively Coupled Plasma–Mass spectrophotometry (ICP-MS) technique to determine the concentration of rare earth and trace elements in the stream sediments.

3.0 Results and Discussion

3.1 Geochemical Composition

Table 1 shows the major oxides component of the Ilubirin stream sediments. The samples are dominated by SiO_2 , which ranges from 72.62-94.81% (Average = 91.7%). The wide range of SiO_2 content may be due to effect of hydraulic sorting and slow deposition. Al_2O_3 ranges from 1.52-5.15% (Average = 2.11%), Fe_2O_3 (Average = 1.84%) MgO (Average = 0.08%); K_2O (Average = 0.37%), TiO_2 (Average = 1.66%), Na_2O (Average = 0.06%), MnO (Average = 0.05) and CaO (Average = 0.08%); this low values may be attributed to chemical destruction under oxidizing conditions during weathering or source-area composition. Lack of Na_2O in the sediments could be attributed to paucity of plagioclase feldspar in the host rock or their loss during weathering.

Figure 2 shows Al_2O_3 correlates positively with all the major oxides except SiO_2 , suggesting hydraulic fractionation and sorting. The negative linear trend between Al_2O_3 and SiO_2 indicates that the major element composition of the stream sediments is controlled largely by the relative amount of quartz and feldspar versus clay minerals. The positive correlation between Al_2O_3 with Fe_2O_3 , MgO and TiO_2 suggests that they occur in clay minerals formed from weathering of ferromagnesian minerals. The positive linear correlation between Al_2O_3 and K_2O suggests that potassium is associated illitic clays. Also, Al_2O_3 correlates positively with P_2O_5 which may be due to association of phosphorus with clays rather than apatite.

Table 1. Major oxides component (Wt %) of the Ilubirin Stream sediments: some ratios and weathering indices

Oxides (Wt. %)	ILUB-1	ILUB-2	ILUB-3	ILUB-4	ILUB-5	ILUB-6	ILUB-7
SiO ₂	94.06	94.63	93.86	94.7	94.21	94.74	94.81
Al ₂ O ₃	1.52	1.78	1.81	1.7	1.65	1.57	1.67
Fe ₂ O ₃	0.87	1.1	0.96	0.83	0.73	0.59	0.94
MnO	0.02	0.02	0.02	0.01	0.01	0.01	0.01
MgO	0.06	0.07	0.06	0.07	0.06	0.06	0.07
K ₂ O	0.03	0.06	0.06	0.06	0.05	0.06	0.05
CaO	0.03	0.04	0.03	0.03	0.03	0.04	0.03
Na ₂ O	0	0	0	0	0	0	0.01
TiO ₂	0.84	0.79	1.14	0.69	0.71	0.58	0.63
P ₂ O ₅	0.04	0.05	0.05	0.04	0.04	0.04	0.04
SiO ₂ /Al ₂ O ₃	61.88	53.16	51.86	55.71	57.10	60.34	56.77
K ₂ O/Al ₂ O ₃	0.02	0.03	0.03	0.04	0.03	0.04	0.03
CaO/MgO	0.50	0.57	0.50	0.43	0.50	0.67	0.43
K ₂ O/Na ₂ O	0	0	0	0	0	0	5.00
Na ₂ O/K ₂ O	0.00	0.00	0.00	0.00	0.00	0.00	0.20
Log(K ₂ O/Na ₂ O)	0	0	0	0	0	0	0.70
Log ((Fe ₂ O ₃ +MgO)/(Na ₂ O+K ₂ O))	1.49	1.29	1.23	1.18	1.20	1.03	1.23
Fe ₂ O ₃ /K ₂ O	29.00	18.33	16.00	13.83	14.60	9.83	18.80
Log SiO ₂ /Al ₂ O ₃	1.79	1.73	1.71	1.75	1.76	1.78	1.75
Log(Fe ₂ O ₃ /K ₂ O)	1.46	1.26	1.20	1.14	1.16	0.99	1.27
PIA	98.03	97.73	98.31	98.2	98.16	97.42	97.59
CIA	96.2	94.68	95.26	94.97	95.38	94.01	94.89
CIW	98.06	97.8	98.37	98.27	98.21	97.52	97.66
MIA	92.41	89.36	90.53	89.94	90.75	88.02	89.77




Fig. 2. Cross-plots of major oxides against Al_2O_3 showing the correlations.

Based on the work by Lindsey (1999) using data from Pettijohn (1963 and 1975), the average lithic arenites plotted in the ferromagnesian potassic sandstones field, but the average greywacke plotted in the sodic sandstone field and average arkoses appeared in the potassic sandstones field. Figure 4 shows a plot in the quartzarenite zone, lack of more plots is due to paucity of Na_2O in the samples.

According to Pettijohn (1963), the lithic arenites are a diverse and poorly defined class. In addition to

abundant rock fragments of widely varying composition, many lithic arenites contain clay matrix with different compositions which can contain higher levels of Fe and Mg. Also, many rock fragments of lithic sandstones are composed of materials that vary greatly in composition. Based on compositional fields for major classes of sandstones (Lindsey, 1999), the studied sediments plotted in the lithic arenite field (Fig. 5).




Figure 3. Ternary diagram of $Na_2O-K_2O-(Fe_2O_3+MgO)$ of the stream sediments, from Blatt et al., 1972.




Figure 4. Chemical classification of the stream sediments based on Pettijohn scheme (1972).




Figure 5. Compositional fields for major classes of sandstones (data from Pettijohn, 1963; 1975): Log $(\text{Fe}_2\text{O}_3+\text{MgO})/(\text{Na}_2\text{O}+\text{K}_2\text{O})$ versus $\log (\text{K}_2\text{O}/\text{Na}_2\text{O})$, adapted from Lindsey, 1999).

Figure 6 shows the samples plotting in the Fe-sand zone. According to Farquhar et al. (2014), the third axis from the Herron (1988) scheme (not shown in Fig. 6), classifies samples by Ca content by dividing

samples into non-calcareous ($\text{Ca} < 4\%$), calcareous ($4\% < \text{Ca} < 15\%$), and carbonate ($\text{Ca} > 15\%$) samples. The stream sediments classified as non-calcareous.




Figure 6. Chemical classification of the stream sediments based on $\log (\text{SiO}_2/\text{Al}_2\text{O}_3)$ vs. $\log (\text{Fe}_2\text{O}_3/\text{K}_2\text{O})$ diagram of Herron (1988).

Based on the study of a reference set, Lindsey (1999) proposed the following guidelines for chemical classification of sandstones:

- 1) quartz arenite: $\log (\text{SiO}_2/\text{Al}_2\text{O}_3) \geq 1.5$
- 2) graywacke: $\log (\text{SiO}_2/\text{Al}_2\text{O}_3) < 1$ and $\log (\text{K}_2\text{O}/\text{Na}_2\text{O}) < 0$

- 3) arkose (includes subarkose): $\log (\text{SiO}_2/\text{Al}_2\text{O}_3) < 1.5$ and $\log (\text{K}_2\text{O}/\text{Na}_2\text{O}) \geq 0$ and $\log ((\text{Fe}_2\text{O}_3+\text{MgO})/(\text{K}_2\text{O}+\text{Na}_2\text{O})) < 0$
- 4) lithic arenite (subgraywacke, includes protoquartzite): $\log (\text{SiO}_2/\text{Al}_2\text{O}_3) < 1.5$ and either $\log (\text{K}_2\text{O}/\text{Na}_2\text{O}) < 0$ or $\log ((\text{Fe}_2\text{O}_3+\text{MgO})/(\text{K}_2\text{O}+\text{Na}_2\text{O})) \geq 0$. If $\log (\text{K}_2\text{O}/\text{Na}_2\text{O}) < 0$, lithic arenite can be confused with graywacke.

The stream sediments falls within the requirements of the first condition and thus classifies as a quartz arenite.

3.3 Maturity

Figure 7 indicates that the stream sediments formed under semi-arid/arid conditions tending towards increasing chemical maturity. The high values given from the ratio of $\text{SiO}_2/\text{Al}_2\text{O}_3$ indicate that all the samples have low degree of clayness. The higher the SiO_2 content the lower the degree of clayness. Potter (1978) stated that maturity of sandstones is reflected by the $\text{SiO}_2/\text{Al}_2\text{O}_3$ index. High ratios indicate mineralogically mature (quartzose, rounded) samples, while low ratios represents chemically immature samples. The $\text{SiO}_2/\text{Al}_2\text{O}_3$ ratio for the Ilubirin stream sediments range between 51.86 and 61.88, which is high; this shows that the samples are mature. The $\text{Na}_2\text{O}/\text{K}_2\text{O}$ and $\text{Fe}_2\text{O}_3/\text{K}_2\text{O}$ ratios can also be used in determining chemical maturity and mineral stability, respectively (Pettijohn et al., 1987; Herron, 1988). The paucity of Na_2O in the samples disallowed the use of the $\text{Na}_2\text{O}/\text{K}_2\text{O}$ ratio. By extrapolation, samples with a low $\text{SiO}_2/\text{Al}_2\text{O}_3$ ratio and a higher $\text{Fe}_2\text{O}_3/\text{K}_2\text{O}$ ratio

should be mineralogically less stable and more prone to reactivity during supercritical CO_2 exposure (Farquhar et al., 2014). The samples studied have a much higher $\text{SiO}_2/\text{Al}_2\text{O}_3$ ratio and low $\text{Fe}_2\text{O}_3/\text{K}_2\text{O}$ ratio, thus, they are mineralogically more stable and less prone to reactivity during supercritical CO_2 exposure. The $\text{Al}_2\text{O}_3/(\text{CaO}+\text{MgO}+\text{Na}_2\text{O}+\text{K}_2\text{O})$ ratio can be used in determining the stability of mobile oxides as proposed by Gill and Yemane (1996). From the positive values obtained (9.8 to 12.7), it shows that there are stable mobile oxides in the stream sediments.

An approach towards assessing detrital mineralogy is to use the Index of Compositional Variability (Cox et al., 1995). The Index of Compositional Variability (ICV) is defined as: $(\text{Fe}_2\text{O}_3+\text{Na}_2\text{O}+\text{CaO}+\text{MgO}+\text{TiO}_2)/\text{Al}_2\text{O}_3$. More matured sandstone with mostly clay minerals displays lower ICV values that are less than 1.0 and such sandstones are derived from cratonic environment (Cox et al., 1995). The results of the Index of Compositional Variability range between 0.81 and 1.21 with an average of 1.03, this indicate that the sands are mineralogically mature.




Figure 7. Chemical maturity of the Ilubirin stream sediment expressed by bivariate plot of SiO_2 versus $\text{Al}_2\text{O}_3 + \text{K}_2\text{O} + \text{Na}_2\text{O}$ (After Suttner & Dutta, 1986).

3.4 Source-area weathering

According to Nesbitt and Young (1982), the evaluation of the degree of chemical weathering of the sediments' source rocks can be determined by calculating the Chemical Index of Alteration (CIA), where $\text{CIA} = \text{molar } (\text{Al}_2\text{O}_3/[\text{Al}_2\text{O}_3+\text{CaO}+\text{Na}_2\text{O}+\text{K}_2\text{O}])$. This index works correctly when Ca, Na, and K decrease as the intensity of weathering increases (Duzgoren-Aydin et al., 2002). The Chemical Index of Weathering (CIW) proposed by Harnois, (1988) is similar to the CIA

except for the exclusion of K_2O in the equation: $\text{CIW} = \text{molar } (\text{Al}_2\text{O}_3/(\text{Al}_2\text{O}_3 + \text{CaO} + \text{Na}_2\text{O}))$. The CIA and CIW are interpreted in similar way with value of 50 for unweathered upper continental crust and roughly 100 for highly weathered materials, with complete removal of alkali and alkaline-earth elements (McLennan et al., 1983; McLennan, 1993; Mongelli et al., 1996). Low CIA values (i.e. 50 or less) also might reflect cool and / or arid conditions (Fedó et al., 1995). The intensity of the chemical weathering can also be estimated using the Plagioclase Index of Alteration

(Fedo et al., 1995); in molecular proportions: $PIA = [(Al_2O_3 - K_2O) / (Al_2O_3 + CaO^* + Na_2O - K_2O)] \times 100$ where CaO^* is the CaO residing only in the silicate fraction. Unweathered plagioclase has PIA value of 50 while Phanerozoic shales have PIA value of 79.

The CIA values for the samples ranged between 94 and 96%, while CIW ranged from 97-98% indicating a high degree of weathering of the source materials. The PIA values ranged from 97-98%, this also indicates high degree of weathering.

Voicu et al. (1997) also proposed the Mineralogical Index of Alteration (MIA) as a weathering parameter calculated as: $MIA = 2 * (CIA - 50)$. MIA values between 0 and 20% are designated as incipient, i.e. just starting; 20-40% (weak); 40-60% (moderate) and 60-100% as intense to extreme degree of weathering. The extreme value of 100% indicates complete weathering of a primary material into its equivalent weathered

product (Voicu and Bardoux, 2002). MIA values for the samples ranged between 88 and 92%, which intense weathering of the source material, this is in agreement with the CIA, CIW PIA.

Figure 8 shows that all the samples plotted at the A end member, which suggests intense chemical weathering and transportation of the sediments. The chemical composition of weathering products in a river basin is expected to exhibit entrenched concepts on mobility of various elements during weathering (Nesbitt et al., 1980; Singh et al., 2005), and therefore to assess the state of chemical and physical weathering (Vital and Statterger, 2000; Singh et al., 2005; Liu et al., 2007). Elemental ratios calculated with respect to Al are used to identify and evaluate the major element mobility. According to Singh et al. (2005), the ratio of the content of element X and Al_2O_3 in rivers divided by the ratio of the same element content of upper continental crust (UCC) gives the elemental ratio.




Fig. 8. Ternary diagram showing the weathering trend of the stream sediments (all in molar proportions); $Al_2O_3 - CaO + Na_2O - K_2O$ (A-CN-K). Fields from Gu et al. (2002).

The elemental ratio refers to the relative enrichment or depletion of the element, i.e., >1 indicates enrichment, <1 indicates depletion, and $=1$ indicates no change in the relative abundance of the element. The stream sediments have, CaO , K_2O , Na_2O , Fe_2O_3 , MgO values less than 1, while SiO_2 , and TiO_2 have values greater than 1. The depletion of highly mobile Na, K and Ca elements is due to leaching during the formation of clay minerals during increased chemical weathering.

The immobile Fe and the less mobile Mg elements were depleted while the immobile Ti was enriched; this suggest that they may are from a felsic source.

Titanium is relatively immobile compared to other elements during sedimentary processes, and hence is a good indicator of source rock composition (McLennan et al., 1993). There is a positive correlation between Ti and Fe (fig. 9); according to Singh, (2009), Fe_2O_3 and

TiO₂ in stream sediments can be expected to be correlate positively due to sorting effects. Linear arrays of data points along lines extending toward the origin in binary Fe₂O₃–TiO₂ plots demonstrate that the elements were immobile, and that they were also hydraulically fractionated in a similar manner (Young,

et al., 2013). The Ilubirin stream sediments have TiO₂/Fe₂O₃ ratio between 0.67 and 1.19, which suggests concentration in the sediments of a heavy mineral phase containing Ti minerals such as, ilmenite and rutile.




Fig. 9. Positive correlation of Fe₂O₃–TiO₂ plots for the stream sediments

3.5 Provenance and tectonic settings

Several authors (Blatt et al., 1980; Bhatia, 1983; Bhatia and Crook, 1986; Roser and Korsch, 1986 & 1988) have related sandstone geochemistry to specific tectonic environment. The discriminant function plot of Roser and Korsch (1988) defined four (4) main provenances: mafic igneous provenance; intermediate igneous provenance; felsic igneous provenance; and quartzose sedimentary provenance (Fig. 10). The stream sediments samples plot in the field P4, quartz-rich sediments of mature continental provenance, associated with a continental passive margin, intracratonic basins, or recycled orogenic provinces. Roser and Korsch (1986) created a tectonic

discrimination diagram using K₂O/Na₂O ratio versus SiO₂ (Fig.11) to determine the tectonic setting of clastic terrigenous sedimentary rocks. The cross plot is used to discriminate between sediments deposited in the Passive Continental Margin (PM), Active Continental Margin (ACM) and the Oceanic Island Arc (OIA). A sample plot in the Passive Margin, lack of more plots is due to the paucity of Na₂O. Figure 12 is another tectonic diagram by Maynard et al., (1982), which also shows the plots in the Passive Margin field, though also affected by the paucity of Na₂O. Figures 13 and 14 are also tectonic discrimination diagrams of the Ilubirin stream sediments indicating the Passive Margin zone.




Fig. 10. Discriminant function diagram using major elements for the provenance signatures of the stream sediments (After Roser & Korsch, 1988).




Fig. 11. Tectonic discrimination plot for the stream sediments (After Roser and Korsch, 1986).




Figure 12. K_2O/Na_2O versus SiO_2/Al_2O_3 ratio – ratio diagram of the stream sediments, after Maynard et al; (1982).
 A1= arc setting and andesitic detritus; A2= evolved arc setting, felsic pluton detritus ACM= Active Continental Margin; PM= Passive Margin.




Fig. 13. Tectonic setting discrimination Plot of Al_2O_3/SiO_2 versus $Fe_2O_3 + MgO$ of the stream sediments. Dashed lines denote the major fields representing various tectonic settings (after Bhatia 1983).




Figure 14. Plot of the major element composition of the stream sediments on the tectonic setting discrimination diagram of Kroonenberg (1994). A: Oceanic island Arc, B: continental island Arc, C: active continental margin, D: passive margin.

4. Conclusions

The sediments can be classified as ferromagnesian potassic, quartz arenites that are non-calcareous. The sediments formed under semi-arid/arid conditions tending towards increasing chemical maturity. The SiO_2/Al_2O_3 index is high, meaning that the samples are mature and there are stable mobile oxides in the sediments. The Index of Compositional Variability values indicate that the sediments are mineralogically mature. The weathering indices indicates a high

degree of weathering of the source materials. The depletion of highly mobile Na, K and Ca elements is due to leaching during the formation of clay minerals during increased chemical weathering. The immobile Fe and the less mobile Mg elements were depleted while the immobile Ti was enriched; this suggest that they may are from a felsic source. The Ilubirin stream sediments TiO_2/Fe_2O_3 ratios suggests concentration in the sediments of a heavy mineral phase containing Ti minerals such as, ilmenite and rutile. There is a

positive correlation between Ti and Fe due to sorting effects and hydraulic fractionation. The tectonic setting is the passive continental margin, while the provenance is the quartz-rich sediments of mature continental provenance, associated with a continental passive margin, intracratonic basins, or recycled orogenic provinces.

References

- I. Armstrong-Altrin, J.S., Lee, Y.I., Verma, S.P., Ramasamy, S. 2004. Geochemistry of sandstones from the upper Miocene kudankulam Formation, southern India: Implication for provenance, weathering and tectonic setting. *Journal of Sedimentary Research*, 74(2): 285 – 297. <http://dx.doi.org/10.1306/082803740285>
- II. Armstrong-Altrin, J.S., Verma, S.P. 2005. Critical evaluation of six tectonic setting discrimination diagrams using geochemical data of Neogene sediments from known tectonic settings: *Sedimentary Geology*, 177(1-2): 115-129. <http://dx.doi.org/10.1016/j.sedgeo.2005.02.004>
- III. Bhatia, M. R. 1983). Plate tectonics and geochemical composition of sandstone. *J. Geol.*, 91: 611-627. <http://dx.doi.org/10.1086/628815>
- IV. Bhatia, M. R. and Crook, K. W. 1986. Trace element characteristics of greywackes and tectonic setting discrimination of sedimentary basins. *Contributions to Mineralogy and Petrology*, 92: p.181-193. <http://dx.doi.org/10.1007/bf00375292>
- V. Bhatt, M.I. and Ghosh, S.K. 2001. Geochemistry of 2.51Ga old Rampur Group pelites, western Himalayas: Implications from their provenance and weathering. *Precambrian Research*, 108, 1-16. [http://dx.doi.org/10.1016/s0301-9268\(00\)00139-x](http://dx.doi.org/10.1016/s0301-9268(00)00139-x)
- VI. Blatt, H., Middleton G. and Murray R. 1972. Origin of sedimentary rocks; Eaglewood cliffs New Jersey Prentice-Hall. p 634. <http://dx.doi.org/10.1017/s0016756800000558>
- VII. Chittleborough, D.J., 1991. Indices of weathering for soils and palaeosols formed on silicate rocks: *Australian Journal of Earth Sciences*, 38, 115–120. <http://dx.doi.org/10.1080/08120099108727959>
- VIII. Cingolani, C.A., Manassero, M., Abre, P. 2003. Composition, provenance, and tectonic setting of Ordovician siliciclastic rocks in the San Rafael block: Southern extension of the Precordillera crustal fragment, Argentina: *Journal of South American Earth Sciences*, 16(1): 91-106. [http://dx.doi.org/10.1016/s0895-9811\(03\)00021-x](http://dx.doi.org/10.1016/s0895-9811(03)00021-x)
- IX. Condie, K.C., 1993. Chemical composition and evolution of the upper continental crust: contrasting results from surface samples and shales. *Chemical Geology*, 104, 1-37. [http://dx.doi.org/10.1016/0009-2541\(93\)90140-e](http://dx.doi.org/10.1016/0009-2541(93)90140-e)
- X. Cox, R., Lowe, D.R., And Cullers, R.L. 1995. The influence of sediment recycling and basement composition on evolution of mudrock chemistry in the southwestern United States: *Geochimica et Cosmochimica Acta*, vol. 59: p. 2919–2940. [http://dx.doi.org/10.1016/0016-7037\(95\)00185-9](http://dx.doi.org/10.1016/0016-7037(95)00185-9)
- XI. Cullers, R.L. and Podkovyrov, V.M. 2000. Geochemistry of the Mesoproterozoic Lakhanda shales in southeastern Yakutia, Russia: Implications for mineralogical and provenance, and recycling. *Precambrian Research*, 104, 77-93. [http://dx.doi.org/10.1016/s0301-9268\(00\)00090-5](http://dx.doi.org/10.1016/s0301-9268(00)00090-5)
- XII. Das, B.K., Haake, B. 2003. Geochemistry of Rewalsar Lake sediments, Lesser Himalaya, India: Implications for source-area weathering, provenance and tectonic setting. *J. Geosci.*, 7: p.299–312. <http://dx.doi.org/10.1007/bf02919560>
- XIII. Dickson, I.N.R. 1985. Interpreting provenance relation from detrital modes of sandstones, in Zuffa G. Gled.), provenance of Arenites: NATO ASI Series (148, D. Reidel Publishing Company, Dordrecht, 333 – 362.
- XIV. Dickinson, W.R., Beard, L.S., Brakenridge, G.R., Erjavec, J.L., Ferguson, R.C., Inman, K.F., Knepp, R.A., Lindberg, F.A. and Ryberg, P.T. 1983. Provenance of North American Phanerozoic sandstones in relation to tectonic setting: *Geological Society of America Bulletin*, 94, 222-235. [http://dx.doi.org/10.1130/0016-7606\(1983\)94%3C222:ponaps%3E2.0.co;2](http://dx.doi.org/10.1130/0016-7606(1983)94%3C222:ponaps%3E2.0.co;2)
- XV. Dickson, W.R., Suczek, C.A. 1979. Plate tectonics and sandstone compositions: American Association of petroleum Geologist, 63: 2164 – 2182.
- XVI. Duzgoren-Aydin, N. S., Aydin, A. and Malpas, J. 2002. Reassessment of chemical weathering indices: case study on pyroclastic rocks of Hong Kong. *Engineering Geology*, 63: p.99–119. [http://dx.doi.org/10.1016/s0013-7952\(01\)00073-4](http://dx.doi.org/10.1016/s0013-7952(01)00073-4)
- XVII. Fairchild, I., Graham, H., Martin, Q., and Maurice, T. 1999. Chemical Analysis of Sedimentary Rocks in: *Techniques in Sedimentology* (ed. T. Maurice), p. 274-354.
- XVIII. Farquhar, S.M., Pearce, J.K., Dawson, G.K.W., Golab, A., Sommacal, S., Kirste, D., Biddle, D., Golding, S.D. 2014. A fresh approach to investigating CO2 storage: Experimental CO2-water-rock interactions in a low-salinity reservoir system, *Chemical Geology*. P. 1-70. <http://dx.doi.org/10.1016/j.chemgeo.2014.10.006>
- XIX. Fedo, C.M., Nesbitt, H. W. & Young, G. M. 1995. Unraveling the effects of potassium metasomatism in sedimentary rocks and paleosols, with implications for paleoweathering conditions and provenance. *Geology*, 23: 10, p.921- 924. [http://dx.doi.org/10.1130/0091-7613\(1995\)023%3C0921:uteopm%3E2.3.co;2](http://dx.doi.org/10.1130/0091-7613(1995)023%3C0921:uteopm%3E2.3.co;2)
- XX. Fedo, C.M., Young, G.M., Nesbitt, H.W. and Hanchar, J.M. 1997. Potassic and sodic metasomatism in the Southern Province of the Canadian Shield: Evidence from the Paleoproterozoic Serpent Formation, Huronian Supergroup. *Canada Precambrian Research*, 84, 17-36. [http://dx.doi.org/10.1016/s0301-9268\(96\)00058-7](http://dx.doi.org/10.1016/s0301-9268(96)00058-7)
- XXI. Folk, R.L. 1974. Petrology of sedimentary rocks. Hemphills Austin Texas. 159pp.
- XXII. Gill, S., Yemane, K. 1996. Implications of a lower Pennsylvanian Ultisol for equatorial Pangean climates and early, oligotrophic, forest ecosystems. *Geology*, Vol. 24: No. 10, p. 905-908. [http://dx.doi.org/10.1130/0091-7613\(1996\)024%3C0905:ioalpu%3E2.3.co;2](http://dx.doi.org/10.1130/0091-7613(1996)024%3C0905:ioalpu%3E2.3.co;2)
- XXIII. Grunsky, E.C., Drew, L.J., Sutphin, D.M. 2009. Process recognition in multi-element soil and stream-sediment geochemical data. *Appl. Geochem.* 24, 1602–1616. <http://dx.doi.org/10.1016/j.apgeochem.2009.04.024>
- XXIV. Gu, X.X., Liu, J.M., Zheng, M.H., Tang, J.X., and Qt, L. 2002. Provenance and Tectonic setting of the Proterozoic turbidites in Hunan, South China: Geochemical Evidence: *Journal of Sedimentary Research*, v. 72, p. 393–407. <http://dx.doi.org/10.1306/081601720393>
- XXV. Halamic, J., Peh, Z., Bukovec, D., Miko, S., Galovic, L. 2001. A factor model of the relationship between stream sediment geochemistry and adjacent drainage basin lithology, Medvednica Mt., Croatia. *Geol. Croat.* 54 (1), 37–51.
- XXVI. Harnois, L. 1988. The C.I.W. index: a new chemical index of weathering. *Sedimentary Geology*, 55: p.319–322. [http://dx.doi.org/10.1016/0037-0738\(88\)90137-6](http://dx.doi.org/10.1016/0037-0738(88)90137-6)
- XXVII. Hendry, J. P. and Trewin, N.H. 1993. Calcite-cemented layers in Deep-marine sandstones: Lower Cretaceous Scapa field, Witch Ground Graben, North Sea. *AAPG*

- Annual Convention, New Orleans, Louisiana. <http://dx.doi.org/10.1306/d9cb58bd-1715-11d7-8645000102c1865d>
- XXVIII. Herron, M. M. 1988. Geochemical classification of terrigenous sands and shales from core or log data. *Journal of Sedimentary Petrology*. 58(5): p.820-829. <http://dx.doi.org/10.1306/212f8e77-2b24-11d7-8648000102c1865d>
- XXIX. Huntsman-Mapilaa, P., Kampunzuc, A.B., Vinkc, B., Ringrosea, S. 2005. Cryptic indicators of provenance from the geochemistry of the Okavango Delta sediments, Botswana. *Sedim. Geol.*, 174: p.123-148. <http://dx.doi.org/10.1016/j.sedgeo.2004.11.001>
- XXX. Igwe, E. O., Amoke, G. U. & Ngwu, C. N. 2013. Provenance and Tectonic Setting of Amasiri Sandstone (Turonian) in Ugep Area, Southern Benue Trough, Nigeria: Evidences from Petrography and Geochemistry. *Global Journal of Science Frontier Research Environment & Earth Science*. Volume 13 Issue 2 Version 1.0: 32-40.
- XXXI. Johnsson, M.J. and Basu, A., 1993. Processes Controlling the Composition of Clastic Sediments. *Geological Society of America*, 284(Special Paper), 342pp. <http://dx.doi.org/10.1130/spe284-pvii>
- XXXII. Jones, H.A. and Hockey R.D. 1964. The Geology of part of Southwestern Nigeria. *Bull. Geol. Surv. Nig.* 31: 101.
- XXXIII. Klemme, H.D. 1975. Geothermal gradient, heat flow and hydrocarbon recovery. In: A.G. Fisher and S. Judson (eds). *Petroleum and global tectonics*. Princeton University Press. p. 251-304.
- XXXIV. Kroonenberg, S.B. 1994: Effects of provenance, sorting and weathering on the geochemistry of fluvial sands from different tectonic and climatic environments: Proceedings of the 29th International Geological Congress, Part A, 69-81.
- XXXV. Kogbe, C.A. 1976. *Geology of Nigeria*. Second revised edition Publ by Rockview Nig. Ltd. 538pp.
- XXXVI. Kovács, J. 2007. Chemical Weathering Intensity of the Late Cenozoic "Red Clay" Deposits in the Carpathian Basin. *Geochemistry International*, Vol. 45: No. 10, pp. 1056-1063. <http://dx.doi.org/10.1134/s0016702907100096>
- XXXVII. Lindsey, D.A. 1999. An Evaluation of Alternative Chemical Classifications of Sandstones. United State Geological Survey Open-File Report, 99-346: 23pp.
- XXXVIII. Liu, Z., Colin, C., Huang, W., Le, K.P., Tong, S., Chen, Z., Trentesaux, A. 2007. Climatic and tectonic controls on weathering in South China and the Indochina Peninsula: clay mineralogical and geochemical investigations from the Pearl, Red, and Mekong drainage basins. *Geochem. Geophys. Geosyst.*, 8: Q05005. [doi:10.1029/2006GC001490](http://dx.doi.org/10.1029/2006GC001490). <http://dx.doi.org/10.1029/2006gc001490>
- XXXIX. Maynard, J.B., Valloni, R. and Yu, H.S. 1982. Composition of modern deep-sea sands from arc-related basins: in Leggett, J.K., eds., *Trench forearc geology: sedimentation and tectonics on modern and ancient active plate margins*. *Geol. Soc. Lond. Spec. Pub.* 10: 551-561
- XL. McLennan, S.M. 1993. Weathering and global denudation. *J. Geol.*, 101: p.295-303. <http://dx.doi.org/10.1086/648222>
- XLI. McLennan, S.M., Taylor, S.R., 1991. Sedimentary rocks and crustal evolution: Tectonic setting and secular trends. *Journal of Geology*, 99, 1-21. <http://dx.doi.org/10.1086/629470>
- XLII. McLennan, S.M., Taylor, S.R., Eriksson, K.A. 1983. Geochemistry of Archean shales from the Pilbara Supergroup, Western Australia. *Geochim. Cosmochim. Acta*, 47: p.1211-1222. [http://dx.doi.org/10.1016/0016-7037\(83\)90063-7](http://dx.doi.org/10.1016/0016-7037(83)90063-7)
- XLIII. McLennan, S.M., Hemming, S., McDaniel, D.K., Hanson, G.N., 1993. Geochemical approaches to sedimentation, provenance and tectonics. *Geol. Soc. Am. Spec. Pap.* 284, 21-40. <http://dx.doi.org/10.1130/spe284-p21>
- XLIV. Milodowski, A.E., and Zalasiewicz, J.A. 1991. Redistribution of rare-earth elements during diagenesis of turbidite / hemipelagite, mudrock sequences of Landoverly age from central Wales. *Geological Society of London, special Publications*, 57, 101-124. *Mining Geol.*, 154 (1): p.65-68. <http://dx.doi.org/10.1144/gsl.sp.1991.057.01.10>
- XLV. Mongelli, G., Cullers, R.L. and Muelheisen, S. 1996. Geochemistry of Late Cretaceous-Oligocene shales from the Varicolori Formation, southern Apennines, Italy: implications for mineralogical, grain-size control and provenance. *Eur. J. Mineral.*, 8: p.733-754. <http://dx.doi.org/10.1127/ejm/8/4/0733>
- XLVI. Nwajide, C.S., and Hoque, M. 1985. Problem of classification and maturity. Evaluation of a diagnostically altered fluvial Sandstone. *Geologic on Nujibouw*, vol.64 p 67-70
- XLVII. Muhs, D.R., Crittenden, R.C., Rosholt, J. N., Bush, C.A., and Stewart, K.C. 1987. Genesis of marine terrace soils, Barbados, West Indies: Evidence from mineralogy and Geochemistry. *Earth Surface Proc. Landforms*, 12: p. 605-618. <http://dx.doi.org/10.1002/esp.3290120604>
- XLVIII. Nesbitt, H.W., and Young, G.M. 1982. Early Proterozoic climates and plate motions inferred from major element chemistry of lutites. *Nature*, 299: 715-717. <http://dx.doi.org/10.1038/299715a0>
- XLIX. Nesbitt, H.W., Mackovics, G., Price, R.C. 1980. Chemical processes affecting alkali sand alkaline Earth during continental weathering. *Geochim. Cosmochim. Acta*, 44: p.1659-1666. [http://dx.doi.org/10.1016/0016-7037\(80\)90218-5](http://dx.doi.org/10.1016/0016-7037(80)90218-5)
- L. Nesbitt, H.W., Young, G.M., McLennan, S.M. and Keays, R.R. 1996. Effects of chemical weathering and sorting on the petrogenesis of siliclastic sediments, with implications for provenance studies. *Journal of Geology*, 104, 525-542. <http://dx.doi.org/10.1086/629850>
- LI. Ogbe, F.G.A. 1972. Stratigraphy of Strata Exposed in Ewekoro Quarry, Southwestern Nigeria. In: *African Geology*, pp 305-322.
- LII. Omatsola, M.E and Adegoke O.S. 1981. Tectonic and Cretaceous stratigraphy of the Dahomey basin. *Journal of Mining Geology*, 154 (1): p.65-68.
- LIII. Omatsola, M.E. and Adegoke, O.S. 1980. Tectonic Evolution of the Dahomey basin [West Africa] and its implication in the opening of the North and South Atlantic. *Broc. 26th Int. Geol. Paris* pp 268.
- LIV. Pettijohn, F. J. 1963. Chemical composition of sandstones—excluding carbonate and volcanic sands, in Fleischer, M., ed., *Data of Geochemistry*, sixth edition, U. S. Geological Survey Professional Paper 440-S: 21 pp.
- LV. Pettijohn, F. J. 1975. *Sedimentary rocks*, third edition: New York, Harper & Row: 628 pp.
- LVI. Pettijohn, F.J., Potter P.E and Siever R. 1972. *Sand and Sandstone*. New York, Springer: 618pp. <http://dx.doi.org/10.1126/science.178.4060.497>
- LVII. Pettijohn, F.J., Potter, P.E., Siever, R., 1987. *Sand and Sandstone*. Springer Science+Business Media, New York, 618 pp. <http://dx.doi.org/10.1007/978-1-4612-1066-5>
- LIX. Potter, P.E., 1978. Petrology and chemistry of modern big river sands. *The Journal of Geology*, 86(4): 423-449. <http://dx.doi.org/10.1086/649711>
- LX. Ratcliffe, K.T., Morton, A.C., Ritcey, D.H. and Evenchick, C.A. 2007. Whole-rock geochemistry and heavy mineral analysis as petroleum exploration tools

- in the Bowser and Sustut basins, British Columbia, Canada. *Bulletin of Canadian Petroleum Geology*, Vol. 55: p. 320–336. <http://dx.doi.org/10.2113/gscpgbull.55.4.320>
- LXI. Reimann, C. and Melezhik, V. 2001. Metallogenic provinces, geochemical provinces and regional geology—what causes large-scale patterns in low density geochemical maps of the C-horizon of podzols in Arctic Europe? *Appl. Geochem.* 16, 963–983. [http://dx.doi.org/10.1016/S0883-2927\(00\)00064-0](http://dx.doi.org/10.1016/S0883-2927(00)00064-0)
- LXII. Roser, B. P. and Korsch, R. J. 1986. Determination of tectonic setting of sandstone-mudstone suites using SiO₂ content and K₂O/Na₂O ratio. *J. Geol.*, 94: p.635-650. <http://dx.doi.org/10.1086/629071>
- LXIII. Roser, B. P. and Korsch, R. J. 1988. Provenance signature of sandstone-mudstone suite determined using discriminant function analysis of major element data. *Chem. Geol.*, 67: p.119-139. [http://dx.doi.org/10.1016/0009-2541\(88\)90010-1](http://dx.doi.org/10.1016/0009-2541(88)90010-1)
- LXIV. Singh, M., Sharma, M., Tobschall, H. L. 2005. Weathering of the Ganga alluvial plain, northern India: implications from fluvial geochemistry of the Gomati River. *Appl. Geochem.*, 20: p.1–21. <http://dx.doi.org/10.1016/j.apgeochem.2004.07.005>
- LXV. Singh, P., 2009. Major, trace and REE geochemistry of the Ganga River sediments: influence of provenance and sedimentary processes. *Chem. Geol.* 266, 242–255. <http://dx.doi.org/10.1016/j.chemgeo.2009.06.013>
- LXVI. Suttner, L.J., Dutta, P.K.. 1986. Alluvial sandstone composition and paleoclimate. L. Framework mineralogy. *Journal of sedimentary petrology*, Vol. 56: p. 329-345. <http://dx.doi.org/10.1306/212f8909-2b24-11d7-8648000102c1865d>
- LXVII. Taylor, S. R., and McLennan, S. M. 1985. *The Continental Crust: Its Composition and Evolution: An Examination of the Geological Record Preserved in Sedimentary Rocks*: Oxford, U.K., Blackwell, 328 pp.
- LXVIII. Vital, H., Statterger, K., 2000. Major and trace elements of stream sediments from the lowermost Amazon River. *Chem. Geol.*, 168: p.151–168. [http://dx.doi.org/10.1016/S0009-2541\(00\)00191-1](http://dx.doi.org/10.1016/S0009-2541(00)00191-1)
- LXIX. Whiteman, A. J. 1982. *Nigeria: Its petroleum geology, resources and potential*, Graham and Trotman, London, Vol 2: 394pp. <http://dx.doi.org/10.1007/978-94-009-7361-9>
- LXX. Wronkiewicz, D.J., and Condie, K.C. 1989. Geochemistry and provenance of sediments from the Pongola Supergroup, South Africa: Evidence for a 3.0-Ga-old continental Craton. *Geochimica et Cosmochimica Acta*, 53: 1537–1549. [http://dx.doi.org/10.1016/0016-7037\(89\)90236-6](http://dx.doi.org/10.1016/0016-7037(89)90236-6)
- LXXI. Wronkiewicz, D.J., Condie, K.C. 1987. Geochemistry of Archean shales from the Witwatersrand Supergroup, South Africa: source-area weathering and provenance. *Geochim. Cosmochim. Acta*, 51: 2401–2416. [http://dx.doi.org/10.1016/0016-7037\(87\)90293-6](http://dx.doi.org/10.1016/0016-7037(87)90293-6)
- LXXII. Young, S.M., Pitawala, A. and Ishiga, H. 2013. Geochemical characteristics of stream sediments, sediment fractions, soils, and basement rocks from the Mahaweli River and its catchment, Sri Lanka. *Chemie der Erde* 73, 357–371. <http://dx.doi.org/10.1016/j.chemer.2012.09.003>
- LXXIII. Zimmermann, U and Bahlburg, H., 2003. Provenance analysis and tectonic setting of the Ordovician clastic deposits in the southern Puna Basin, NW Argentina. *Sedimentology*, 50, 1079–1104. <http://dx.doi.org/10.1046/j.1365-3091.2003.00595.x>